

## Geometry

---

### Objectives

- Introduce the elements of geometry
  - Scalars
  - Vectors
  - Points
- Develop mathematical operations among them in a coordinate-free manner
- Define basic primitives
  - Line segments
  - Polygons

## Basic Elements

---

- Geometry is the study of the relationships among objects in an n-dimensional space
  - In computer graphics, we are interested in objects that exist in three dimensions
- Want a minimum set of primitives from which we can build more sophisticated objects
- We will need three basic elements
  - Scalars
  - Vectors
  - Points

## Coordinate-Free Geometry

---

- When we learned simple geometry, most of us started with a Cartesian approach
  - Points were at locations in space  $\mathbf{p}=(x,y,z)$
  - We derived results by algebraic manipulations involving these coordinates
- This approach was nonphysical
  - Physically, points exist regardless of the location of an arbitrary coordinate system
  - Most geometric results are independent of the coordinate system
  - Example Euclidean geometry: two triangles are identical if two corresponding sides and the angle between them are identical


## Scalars

---

- Need three basic elements in geometry
  - Scalars, Vectors, Points
- Scalars can be defined as members of sets which can be combined by two operations (addition and multiplication) obeying some fundamental axioms (associativity, commutivity, inverses)
- Examples include the real and complex number systems under the ordinary rules with which we are familiar
- Scalars alone have no geometric properties

## Vectors

- Physical definition: a vector is a quantity with two attributes
  - Direction
  - Magnitude
- Examples include
  - Force
  - Velocity
  - Directed line segments
    - Most important example for graphics
    - Can map to other types




Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 5

## Vector Operations

- Every vector has an inverse
  - Same magnitude but points in opposite direction
- Every vector can be multiplied by a scalar
  - Zero magnitude, undefined orientation
- The sum of any two vectors is a vector
  - Use head-to-tail axiom



Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 6


## Linear Vector Spaces

- Mathematical system for manipulating vectors
  - Operations
    - Scalar-vector multiplication  $u = \alpha v$
    - Vector-vector addition:  $w = u + v$
  - Expressions such as  $v = u + 2w - 3r$
- Make sense in a vector space

Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 7

## Vectors Lack Position


- These vectors are identical
    - Same length and magnitude
- 
- Vectors spaces insufficient for geometry
    - Need points

Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 8

## Points

- Location in space
- Operations allowed between points and vectors
  - Point-point subtraction yields a vector
  - Equivalent to point-vector addition



Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 9

## Affine Spaces


- Point + a vector space
- Operations
  - Vector-vector addition
  - Scalar-vector multiplication
  - Point-vector addition
  - Scalar-scalar operations
- For any point define
  - $1 \cdot P = P$
  - $0 \cdot P = \mathbf{0}$  (zero vector)

Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 10

## Lines

- Consider all points of the form
  - $P(\alpha) = P_0 + \alpha \mathbf{d}$
  - Set of all points that pass through  $P_0$  in the direction of the vector  $\mathbf{d}$



Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 11

## Parametric Form

- This form is known as the parametric form of the line
  - More robust and general than other forms
  - Extends to curves and surfaces
- Two-dimensional forms
  - Explicit:  $y = mx + h$
  - Implicit:  $ax + by + c = 0$
  - Parametric:
    - $x(\alpha) = \alpha x_0 + (1-\alpha)x_1$
    - $y(\alpha) = \alpha y_0 + (1-\alpha)y_1$

Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 12

## Rays and Line Segments


- If  $\alpha \geq 0$ , then  $P(\alpha)$  is the *ray* leaving  $P_0$  in the direction  $\mathbf{d}$

If we use two points to define  $\mathbf{v}$ , then

$$P(\alpha) = Q + \alpha(R - Q) = Q + \alpha\mathbf{v}$$

$$= \alpha R + (1 - \alpha)Q$$

For  $0 \leq \alpha \leq 1$  we get all the points on the *line segment* joining R and Q




Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005


KENT STATE 13

## Convexity

- An object is *convex* iff for any two points in the object all points on the line segment between these points are also in the object



convex



not convex

Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 14

## Affine Sums

- Consider the “sum”

$$P = \alpha_1 P_1 + \alpha_2 P_2 + \dots + \alpha_n P_n$$

Can show by induction that this sum makes sense iff

$$\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$$

in which case we have the *affine sum* of the points  $P_1, P_2, \dots, P_n$


- If, in addition,  $\alpha_i \geq 0$ , we have the *convex hull* of  $P_1, P_2, \dots, P_n$

Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 15

## Convex Hull

- Smallest convex object containing  $P_1, P_2, \dots, P_n$
- Formed by “shrink wrapping” points




Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 16

## Curves and Surfaces

- Curves are one parameter entities of the form  $P(\alpha)$  where the function is nonlinear
- Surfaces are formed from two-parameter functions  $P(\alpha, \beta)$ 
  - Linear functions give planes and polygons




Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 17

## Planes


- A plane can be defined by a point and two vectors or by three points



Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 18

## Triangles




Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 19

## Normals

- Every plane has a vector  $n$  normal (perpendicular, orthogonal) to it
- From point-two vector form  $T(\alpha, \beta) = P_0 + \alpha u + \beta v$ , we know we can use the cross product to find  $n = u \times v$  and the equivalent form for the plane  $(P(\alpha) - P_0) \cdot n = 0$



Angel: Interactive Computer Graphics 4E © Addison-Wesley 2005

KENT STATE 20