

Predicates and Quantifiers

Section 1.4

Section Summary

- Predicates
- Variables
- Quantifiers
 - Universal Quantifier
 - Existential Quantifier
- Negating Quantifiers
 - De Morgan's Laws for Quantifiers
- Translating English to Logic

Propositional Logic Not Enough

- If we have:
 - “All men are mortal.”
 - “Socrates is a man.”
- Does it follow that “Socrates is mortal?”
- Can't be represented in propositional logic. Need a language that talks about objects, their properties, and their relations.
- Later we'll see how to draw inferences.

Introducing Predicate Logic

- Predicate logic uses the following new features:
 - Variables: x, y, z
 - Predicates: $P(x), M(x)$
 - Quantifiers (*to be covered in a few slides*):
- *Propositional functions* are a generalization of propositions.
 - They contain variables and a predicate, e.g., $P(x)$
 - Variables can be replaced by elements from their *domain*.

Propositional Functions

- Propositional functions become propositions (and have truth values) when their variables are each replaced by a value from the *domain* (or *bound* by a quantifier, as we will see later).
- The statement $P(x)$ is said to be the value of the propositional function P at x .
- For example, let $P(x)$ denote “ $x > 0$ ” and the domain be the integers. Then:
 - $P(-3)$ is false.
 - $P(0)$ is false.
 - $P(3)$ is true.
- Often the domain is denoted by U . So in this example U is the integers.

Examples of Propositional Functions

- Let “ $x + y = z$ ” be denoted by $R(x, y, z)$ and U (for all three variables) be the integers. Find these truth values:

$R(2, -1, 5)$

Solution: F

$R(3, 4, 7)$

Solution: T

$R(x, 3, z)$

Solution: Not a Proposition

- Now let “ $x - y = z$ ” be denoted by $Q(x, y, z)$, with U as the integers. Find these truth values:

$Q(2, -1, 3)$

Solution: T

$Q(3, 4, 7)$

Solution: F

$Q(x, 3, z)$

Solution: Not a Proposition

Compound Expressions

- Connectives from propositional logic carry over to predicate logic.
- If $P(x)$ denotes “ $x > 0$,” find these truth values:
 - $P(3) \vee P(-1)$ **Solution:** T
 - $P(3) \wedge P(-1)$ **Solution:** F
 - $P(3) \rightarrow P(-1)$ **Solution:** F
 - $P(3) \rightarrow P(-1)$ **Solution:** T
- Expressions with variables are not propositions and therefore do not have truth values. For example,
 - $P(3) \wedge P(y)$
 - $P(x) \rightarrow P(y)$
- When used with quantifiers (to be introduced next), these expressions (propositional functions) become propositions.



Charles Peirce (1839-1914)

Quantifiers

- We need *quantifiers* to express the meaning of English words including *all* and *some*:
 - “All men are Mortal.”
 - “Some cats do not have fur.”
- The two most important quantifiers are:
 - *Universal Quantifier*, “For all,” symbol: \forall
 - *Existential Quantifier*, “There exists,” symbol: \exists
- We write as in $\forall x P(x)$ and $\exists x P(x)$.
- $\forall x P(x)$ asserts $P(x)$ is true for every x in the *domain*.
- $\exists x P(x)$ asserts $P(x)$ is true for some x in the *domain*.
- The quantifiers are said to bind the variable x in these expressions.

Universal Quantifier

- $\forall x P(x)$ is read as “For all x , $P(x)$ ” or “For every x , $P(x)$ ”

Examples:

- 1) If $P(x)$ denotes “ $x > 0$ ” and U is the integers, then $\forall x P(x)$ is false.
- 2) If $P(x)$ denotes “ $x > 0$ ” and U is the positive integers, then $\forall x P(x)$ is true.
- 3) If $P(x)$ denotes “ x is even” and U is the integers, then $\forall x P(x)$ is false.

Existential Quantifier

- $\exists x P(x)$ is read as “For some x , $P(x)$ ”, or as “There is an x such that $P(x)$,” or “For at least one x , $P(x)$.”

Examples:

1. If $P(x)$ denotes “ $x > 0$ ” and U is the integers, then $\exists x P(x)$ is true. It is also true if U is the positive integers.
2. If $P(x)$ denotes “ $x < 0$ ” and U is the positive integers, then $\exists x P(x)$ is false.
3. If $P(x)$ denotes “ x is even” and U is the integers, then $\exists x P(x)$ is true.

Uniqueness Quantifier (*optional*)

- $\exists!x P(x)$ means that $P(x)$ is true for one and only one x in the universe of discourse.
- This is commonly expressed in English in the following equivalent ways:
 - “There is a unique x such that $P(x)$.”
 - “There is one and only one x such that $P(x)$ ”
- Examples:
 1. If $P(x)$ denotes “ $x + 1 = 0$ ” and U is the integers, then $\exists!x P(x)$ is true.
 2. But if $P(x)$ denotes “ $x > 0$,” then $\exists!x P(x)$ is false.
- The uniqueness quantifier is not really needed as the restriction that there is a unique x such that $P(x)$ can be expressed as:

$$\exists x (P(x) \wedge \forall y (P(y) \rightarrow y=x))$$

Thinking about Quantifiers

- When the domain of discourse is finite, we can think of quantification as looping through the elements of the domain.
- To evaluate $\forall x P(x)$ loop through all x in the domain.
 - If at every step $P(x)$ is true, then $\forall x P(x)$ is true.
 - If at a step $P(x)$ is false, then $\forall x P(x)$ is false and the loop terminates.
- To evaluate $\exists x P(x)$ loop through all x in the domain.
 - If at some step, $P(x)$ is true, then $\exists x P(x)$ is true and the loop terminates.
 - If the loop ends without finding an x for which $P(x)$ is true, then $\exists x P(x)$ is false.
- Even if the domains are infinite, we can still think of the quantifiers this fashion, but the loops will not terminate in some cases.

Properties of Quantifiers

- The truth value of $\exists x P(x)$ and $\forall x P(x)$ depend on both the propositional function $P(x)$ and on the domain U .
- **Examples:**
 1. If U is the positive integers and $P(x)$ is the statement “ $x < 2$ ”, then $\exists x P(x)$ is true, but $\forall x P(x)$ is false.
 2. If U is the negative integers and $P(x)$ is the statement “ $x < 2$ ”, then both $\exists x P(x)$ and $\forall x P(x)$ are true.
 3. If U consists of 3, 4, and 5, and $P(x)$ is the statement “ $x > 2$ ”, then both $\exists x P(x)$ and $\forall x P(x)$ are true. But if $P(x)$ is the statement “ $x < 2$ ”, then both $\exists x P(x)$ and $\forall x P(x)$ are false.

Precedence of Quantifiers

- The quantifiers \forall and \exists have higher precedence than all the logical operators.
- For example, $\forall x P(x) \vee Q(x)$ means $(\forall x P(x)) \vee Q(x)$
- $\forall x (P(x) \vee Q(x))$ means something different.
- Unfortunately, often people write $\forall x P(x) \vee Q(x)$ when they mean $\forall x (P(x) \vee Q(x))$.

Translating from English to Logic

Example 1: Translate the following sentence into predicate logic: “Every student in this class has taken a course in Java.”

Solution:

First decide on the domain U .

Solution 1: If U is all students in this class, define a propositional function $J(x)$ denoting “ x has taken a course in Java” and translate as $\forall x J(x)$.

Solution 2: But if U is all people, also define a propositional function $S(x)$ denoting “ x is a student in this class” and translate as $\forall x (S(x) \rightarrow J(x))$.

$\forall x (S(x) \wedge J(x))$ is not correct. What does it mean?

Translating from English to Logic

Example 2: Translate the following sentence into predicate logic: “Some student in this class has taken a course in Java.”

Solution:

First decide on the domain U .

Solution 1: If U is all students in this class, translate as

$$\exists x J(x)$$

Solution 1: But if U is all people, then translate as

$$\exists x (S(x) \wedge J(x))$$

$\exists x (S(x) \rightarrow J(x))$ is not correct. What does it mean?

Returning to the Socrates Example

- Introduce the propositional functions $Man(x)$ denoting “ x is a man” and $Mortal(x)$ denoting “ x is mortal.” Specify the domain as all people.
- The two premises are: $\forall x Man(x) \rightarrow Mortal(x)$
 $Man(Socrates)$
- The conclusion is: $Mortal(Socrates)$
- Later we will show how to prove that the conclusion follows from the premises.

Equivalences in Predicate Logic

- Statements involving predicates and quantifiers are *logically equivalent* if and only if they have the same truth value
 - for every predicate substituted into these statements and
 - for every domain of discourse used for the variables in the expressions.
- The notation $S \equiv T$ indicates that S and T are logically equivalent.
- **Example:** $\forall x \neg\neg S(x) \equiv \forall x S(x)$

Thinking about Quantifiers as Conjunctions and Disjunctions

- If the domain is finite, a universally quantified proposition is equivalent to a conjunction of propositions without quantifiers and an existentially quantified proposition is equivalent to a disjunction of propositions without quantifiers.
- If U consists of the integers 1, 2, and 3:

$$\forall x P(x) \equiv P(1) \wedge P(2) \wedge P(3)$$

$$\exists x P(x) \equiv P(1) \vee P(2) \vee P(3)$$

- Even if the domains are infinite, you can still think of the quantifiers in this fashion, but the equivalent expressions without quantifiers will be infinitely long.

Negating Quantified Expressions

- Consider $\forall x J(x)$

“Every student in your class has taken a course in Java.”

Here $J(x)$ is “x has taken a course in Java” and the domain is students in your class.

- Negating the original statement gives “It is not the case that every student in your class has taken Java.” This implies that “There is a student in your class who has not taken Java.”

Symbolically $\neg \forall x J(x)$ and $\exists x \neg J(x)$ are equivalent

Negating Quantified Expressions (continued)

- Now Consider $\exists x J(x)$

“There is a student in this class who has taken a course in Java.”

Where $J(x)$ is “x has taken a course in Java.”

- Negating the original statement gives “It is not the case that there is a student in this class who has taken Java.” This implies that “Every student in this class has not taken Java”

Symbolically $\neg \exists x J(x)$ and $\forall x \neg J(x)$ are equivalent

De Morgan's Laws for Quantifiers

- The rules for negating quantifiers are:

<i>Negation</i>	<i>Equivalent Statement</i>	<i>When Is Negation True?</i>	<i>When False?</i>
$\neg\exists x P(x)$	$\forall x\neg P(x)$	For every x , $P(x)$ is false.	There is an x for which $P(x)$ is true.
$\neg\forall x P(x)$	$\exists x\neg P(x)$	There is an x for which $P(x)$ is false.	$P(x)$ is true for every x .

- The reasoning in the table shows that:

$$\neg\forall x P(x) \equiv \exists x\neg P(x)$$

$$\neg\exists x P(x) \equiv \forall x\neg P(x)$$

- These are important. You will use these.

Translation from English to Logic

Examples:

1. “Some student in this class has visited Mexico.”

Solution: Let $M(x)$ denote “ x has visited Mexico” and $S(x)$ denote “ x is a student in this class,” and U be all people.

$$\exists x (S(x) \wedge M(x))$$

2. “Every student in this class has visited Canada or Mexico.”

Solution: Add $C(x)$ denoting “ x has visited Canada.”

$$\forall x (S(x) \rightarrow (M(x) \vee C(x)))$$